

FTUV-00-1204A
 IFIC/00-78
 hep-th/0012019

Evaporation of charged black holes near extremality*

A. Fabbri^{a†}, D. J. Navarro^{b‡} and J. Navarro-Salas^{b§}

a) Dipartimento di Fisica dell'Università di Bologna and INFN sezione di Bologna,
 Via Irnerio 46, 40126 Bologna, Italy.

b) Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC.
 Facultad de Física, Universidad de Valencia, Burjassot-46100, Valencia, Spain.

The quantum evolution of black holes including back reaction effects is an open problem. One can simplify it by considering the scattering of low-energy neutral particles off extremal Reissner-Nordström black holes [1]. If, in addition, we also consider spherically reduced configurations, the theory can be described, in a region very close to the horizon, by a solvable effective model. This model is just the Jackiw-Teitelboim model [2] (for details see [3])

$$I = \int d^2x \sqrt{-g} \left[\left(R + \frac{4}{l^2 q^3} \right) \tilde{\phi} - \frac{1}{2} \sum_i^N |\nabla f_i|^2 \right], \quad (1)$$

where the two-dimensional fields of (1) are related to the four-dimensional metric by the expression $ds_{(4)}^2 = \frac{2l}{r_0} ds_{(2)}^2 + (r_0^2 + 4l^2 \tilde{\phi}) d\Omega^2$, $r_0 = lq$ is the extremal radius, and $l^2 = G$ is Newton's constant. The fields f_i represent null matter. To analyze back reaction effects we have to add to the classical action (1) the Polyakov-Liouville term

$$I_{PL} = -\frac{N\hbar}{96\pi} \int d^2x \sqrt{-g} R \square^{-1} R + \frac{N\hbar}{12\pi} \int d^2x \sqrt{-g} \lambda^2. \quad (2)$$

The above expression contains a cosmological term with $\lambda^2 = l^{-2} q^{-3}$ to ensure that the extremal geometry remains an exact solution of the one-loop theory. The equations of motion derived from $I + I_{PL}$ in conformal gauge $ds^2 = -e^{2\rho} dx^+ dx^-$ are

$$2\partial_+ \partial_- \rho + \lambda^2 e^{2\rho} = 0, \quad \partial_+ \partial_- \tilde{\phi} + \lambda^2 \tilde{\phi} e^{2\rho} = 0, \quad \partial_+ \partial_- f_i = 0, \quad (3)$$

$$-2\partial_\pm^2 \tilde{\phi} + 4\partial_\pm \rho \partial_\pm \tilde{\phi} = T_{\pm\pm}^f - \frac{N\hbar}{12\pi} t_\pm - \frac{N\hbar}{12\pi} ((\partial_\pm \rho)^2 - \partial_\pm^2 \rho). \quad (4)$$

*Talk given at the Ninth Marcel Grossmann Meeting.

†fabbria@bo.infn.it

‡dnavarro@ific.uv.es

§jnavarro@ific.uv.es

The first equation implies that the two-dimensional metric has a negative constant curvature. To find solutions to the above equations it is convenient to choose the following form of the metric

$$ds^2 = -\frac{2l^2 q^3 dx^+ dx^-}{(x^- - x^+)^2}. \quad (5)$$

This way only the $t_{\pm}$ terms survive in the quantum correction to the equations of motion. However it is the choice of those functions $t_{\pm}$ the crucial point to get the physical solutions. If u and v are the null coordinates associated to the Eddington-Finkelstein coordinates of the Reissner-Nordström metric, the absence of ingoing and outgoing fluxes at extremality (where $u = x^-$ and $v = x^+$) is given by the conditions $t_{vv}(v) = 0 = t_{uu}(u)$. However, once we send matter the relation $v = v(x^+)$ is not the identity although the coordinate u remains equal to x^- . This means that in the general situation we have that $t_-(x^-) = 0$ and $t_+(x^+) = \frac{1}{2}\{v, x^+\}$. The incoming quantum flux will be given by $\langle T_{vv}^f(v) \rangle = -\frac{N\hbar}{12\pi} [(\partial_v \rho)^2 - \partial_v^2 \rho]$, which turns out to be

$$\langle T_{vv}^f(v) \rangle = -\frac{N\hbar}{12\pi} \{v, x^+\} \left(\frac{dx^+}{dv} \right)^2. \quad (6)$$

The equations of motion can be solved immediately in the Vadya-type metric

$$ds^2 = -\left(\frac{2\tilde{x}^2}{l^2 q^3} - l\tilde{m}(v) \right) dv^2 + 2d\tilde{x}dv, \quad \tilde{\phi} = \frac{\tilde{x}}{l}, \quad (7)$$

where $\frac{d\tilde{m}(v)}{dv} = \langle T_{vv}^f(v) \rangle + T_{vv}^f$. The problem is to find the relation between x^+ and v coordinates. Defining $F(x^+)$ as $F = lq^3 \frac{dx^+}{dv}$ and jumping from (7) to the conformal gauge we find that

$$\tilde{\phi} = \frac{F(x^+)}{x^- - x^+} + \frac{1}{2} F'(x^+). \quad (8)$$

The equations of motion, in conformal gauge, imply the following differential equation for F

$$F''' = \frac{N\hbar}{24\pi} \left(-\frac{F''}{F} + \frac{1}{2} \left(\frac{F'}{F} \right)^2 \right) - T_{++}^f(x^+). \quad (9)$$

In terms of the mass function the evolution equation is then

$$\partial_v \tilde{m}(v) = -\frac{N\hbar}{24\pi l q^3} \tilde{m}(v) + T_{vv}^f, \quad (10)$$

and, at late times, $m(v)$ is given by $\tilde{m}(v) = \Delta m e^{-\frac{N\hbar}{24\pi l q^3}(v-v_0)}$. This implies that as time passes the evaporating black hole approaches the extremal configuration. Moreover the equation (10) suggests that the incoming (negative) quantum flux contains the information of the classical stress tensor T_{vv}^f .

References

- [1] A. Strominger and S.P. Trivedi, *Phys. Rev.* **D48**, 5778 (1993); D.A. Lowe and M. O’Loughlin, *Phys. Rev.* **D48**, 3735 (1993).
- [2] R. Jackiw, in “Quantum Theory of Gravity”, edited by S.M. Christensen (Hilger, Bristol, 1984), p. 403; C. Teitelboim, in op. cit., p. 327.
- [3] A. Fabbri, D. J. Navarro and J. Navarro-Salas, *Phys. Rev. Lett.* **85**, 2434 (2000); *Quantum evolution of near-extremal Reissner-Nordström black holes*, hep-th/0006035 (to appear in *Nucl. Phys.* **B**).